

1 Theriogenology
(2008) 69 :870–876



Effects of catecholamines on ovary morphology, blood concentrations of estradiol-17b, progesterone, zinc, triglycerides and rate of ovulation in domestic hens

T.A. Ebeid ^{a,*}, Y.Z. Eid ^a, E.A. El-Abd ^b, M.M. El-Habbak ^a

^a Department of Poultry Production, Faculty of Agriculture, Kafrelsheikh University, 33516 Kafr El Sheikh, Egypt

^b Animal Production Research Institute, Ministry of Agriculture, Dokki, Giza, Egypt

Abstract

The present study is an attempt to shed more light on the role of epinephrine (EP) and norepinephrine (NE) in regulating ovarian follicular development, folliculogenesis and ovulation in laying hens. Sixty Egyptian local cross females (Mandarah), 50 weeks old, were individually housed and equally divided into three treatments: control (saline, 0.9% NaCl), EP (0.15 mg epinephrine/hen/day) and NE (0.75 mg norepinephrine/hen/day) (n = 20). Animals were injected intramuscularly once a day for 15 successive days. At the end of the experimental period, 10 females from each treatment were randomly chosen, weighed and killed by decapitation. Ovaries and oviducts and ovarian follicles were examined. Plasma concentrations of estradiol-17b, progesterone, zinc and triglyceride were determined. Results indicated that the ovaries of NE- and EP-treated hens were more developed than those of control hens being heavier and containing more yellow yolk-filled follicles. EP or NE significantly increased the ovulation rate and plasma concentrations of estradiol-17b, progesterone, zinc and triglyceride compared with control treatment. It could be concluded that catecholamines may have a part in promoting ovarian follicular development and in stimulating ovulation in laying hens at the end of their reproductive lives.



2

Tree Genetics & Genomes
(2008) 4:611–623



Phylogeography of *Larix sukaczewii* Dyl. and *Larix Sibirica* L. inferred from nucleotide variation of nuclear genes

Neyton H. T. Araki & Ismael A. Khatib & Kariyawasam K. G. U. Hemamali & Nobuyuki Inomata & Xiao-Ru Wang & Alfred E. Szmidt

Department of Genetics, Faculty of Agriculture,
Kafri El-Sheikh University,
Kafri El-Sheikh, Egypt

Abstract

We investigated phylogeography of *Larix sukaczewii* and *Larix sibirica* using nucleotide variation at three following nuclear gene regions: 5.8 S rDNA including two internal transcribed spacers (ITS), cinnamyl alcohol dehydrogenase (CAD), and phytochrome-O (PHYO). We also included sequences of the 4-coumarate: coenzyme A ligase (4CL) gene region obtained in our recent study. CAD and PHYO showed very low nucleotide variation, but ITS and 4CL had levels of variation similar to those reported for other conifers. Pleistocene refugia have been hypothesized to exist in the Southern Urals and South Central Siberia, where four out of nine of the investigated populations occur. We found moderate to high levels of population differentiation ($F_{ST}=0.115-0.531$) in some pairwise comparisons suggesting limited gene flow and independent evolution of some refugial populations. In *L. sukaczewii*, low levels of differentiation were found among populations from areas glaciated during the Pleistocene, indicating their recent origin. Our results also suggest these populations were created by migrants from multiple, genetically distinct refugia. Furthermore, some haplotypes observed in populations from previously glaciated areas were not found in putative refugial populations, suggesting these populations might have contributed little to the extant populations created after the Last Glacial Maximum. Some authors regard *L. sukaczewii* and *L. sibirica* as a single species, while others consider them as separate species. The observed conspicuous differences in haplotype composition and distribution between *L. sukaczewii* and *L. sibirica*, together with high values of F_{ST} between populations of the two species, appear to support the latter classification.



٣

Genes Genet. Syst.
(2008) 83 : 55–66



Phylogeography of Eurasian *Larix* species inferred from nucleotide variation in two nuclear genes

Ismael A. Khatab^{1,3}, Hiroko Ishiyama¹, Nobuyuki Inomata¹,
Xiao-Ru Wang² and Alfred E. Szmidt^{1*}

¹Department of Biology, Faculty of Science, Kyushu University, Fukuoka 812-8581, Japan

²Department of Ecology and Environmental Science, Umeå University, 901 87 Umeå, Sweden

³Department of Genetics, Faculty of Agriculture, Kafar El-Sheikh University, Egypt

Abstract

Larch (*Larix* Mill.) is one of the most widely distributed tree genera in Eurasia. To determine population structure and to verify classification of five species and three varieties of the Eurasian *Larix* species, we investigated levels and patterns of nucleotide variation of two nuclear gene regions: the 4-coumarate coenzyme A ligase (4CL) and the coumarate 3-hydroxylase (C3H). In the 4CL region nucleotide diversity at silent sites (π_{sil}) varied between 0.0020 in *L. Gmelinii* to 0.0116 in *L. gmelinii* var. *japonica* and in the C3H region between 0.0019 in *L. kaempferi* to 0.0066 in *L. gmelinii* var. *japonica*. In both gene regions statistically significant population differentiation (F_{ST}) was detected among adjacent refugial populations of some species suggesting limited gene flow and/or long time isolation of some refugial populations. On the other hand, populations of *L. sukaczewii* from northwestern Russia, which was glaciated 20,000 years ago showed no differentiation. This result is consistent with recent postglacial origin of these populations. Haplotype composition of some of the investigated Eurasian *Larix* species suggested that they are considerably diverged. Some haplotypes were unique to individual species. Our results indicate that more intensive sampling especially from known refugial regions is necessary for inferring correct classification of Eurasian *Larix* species and inferring their postglacial migration.

4 Acta Biologica Szegediensis
(2008) 52 : 27-31



In vitro propagation of two triploid hybrids of watermelon through adventitious shoot organogenesis and shoot tip culture

Tarek A. Shalaby^{1*}, Soliman A. Omran², Yousry A. Baioumi¹

¹Horticulture Department, Faculty of Agriculture, Kafrelsheikh University, Egypt, ²Horticulture Research Institute, Agriculture Research Center, Giza, Egypt

Abstract

In vitro propagation protocol for two triploid hybrids of watermelon using cotyledon explants and shoot tips was achieved. Five benzyladenine (BA) concentrations were tested using cotyledon and shoot tip explants. Cotyledon explants and shoot tips from 6 and 15-20 days aseptically germinated were cultured on Murashige and Skoog medium (MS) containing test concentration of benzyladenine (2.22, 4.44, 10, 24.61 and 44.4 μM). Adventitious shoot organogenesis was initiated in all induction media and the differences among BA concentration were significant. MS medium containing 4.44, 10 and 24.61 μM BA showed the highest percentage of explants with shoots. The stimulation of axillary-bud development from excised shoot tips by a high cytokinin (BA) was observed. Axillary shoots were obtained from shoot tips of triploid watermelon and the multiplication rate ranged from 2 to 5.6 plants dependence on benzyladenine concentration and genotype. Obtained data showed that variation in regeneration rate was demonstrated. Shoots were excised and elongated in MS medium without hormones. The elongated shoots were rooted in MS medium containing 0.1 μM -naphthalene acetic acid (NAA). Rooted plants were successfully acclimatized and gradually hardened-off to green-house conditions and subsequently established in soil with a survival rate of 80%.

5

Environmental and Experimental Botany
(2008) 63 : 297–304

Effect of carbon dioxide on antioxidant enzymes and ginsenoside production in root suspension cultures panax ginseng

Mohammad Babar Ali a,., Yaser Hassan Dewir b, Eun-Joo Hahnc, Kee-Yoeup Paek c,1

a Metabolic Regulation Laboratory, Food Biotechnology Division, National Food Research Institute, Ibaraki, Tsukuba, Kannondai 305-8642, Japan b Department of Horticulture, Faculty of Agriculture, Kafr El-Sheikh University, Kafr El-Sheikh 33516, Egypt c Research Center for the Development of Advanced Horticultural Technology, Chungbuk National University, Cheong-ju 361-763, Republic of Korea

Abstract

The aim of this study was to investigate the effect of CO₂ at various concentrations (1, 2.5 and 5%) on antioxidant enzymes and ginsenoside accumulation in *Panax ginseng* roots in 5 l airlift bioreactors (working volume 4 l). One and 2.5% CO₂ was beneficial for root biomass accumulation, but 5% CO₂ decreased the biomass. Ginsenoside concentration decreased with increasing concentration of CO₂. No significant difference was observed in the malondialdehyde (MDA) content and lipoxygenase (LOX) activity between respective controls and CO₂ treated roots. Antioxidant enzymes such as ascorbate peroxidase (APX), monodehydroascorbate reductase (MDHAR), glutathione reductase (GR), catalase (CAT), guaiacol peroxidase (G-POD) including reduced ascorbate and total glutathione were induced in CO₂ exposed roots which emphasized the protective role of antioxidants against CO₂ induced stress. Superoxide dismutase activity (SOD) which was induced after 15 days was significantly inhibited after 45 days. Glutathione-S-transferase (GST) and glutathione peroxidase (GPX) activities also increased when the roots were subjected to 1 and 2.5% CO₂ compared to the respective controls but not at 5%. A higher reduced ascorbate to oxidized (ASC/DHA) ratio in CO₂ treated root indicates the plant's ability to tolerate CO₂ stress. These observations suggest that an increase in antioxidant enzymes may affect a defense response to the cellular damage induced by CO₂. Probably, this increase could not stop the deleterious effects of CO₂ concentration on ginsenoside concentration, but reduced stress severity and thereby allowing root growth to occur.



6

Plant Growth Regul
(2008) 54:157–164



Kinetics of nutrient utilization and photosynthetic enzyme activities during floral versus vegetative differentiation of *Spathiphyllum* in air-lift bioreactor cultures

Yaser Hassan Dewir Debasis Chakrabarty Eun Joo Hahn Subodh Kumar
Datta Kee Yoeup Paek


Department of Horticulture, Faculty of Agriculture, Kafr El-Sheikh University,
Kafr El-Sheikh 33516, Egypt

Abstract

The present study reports on the kinetics of nutrient utilization during in vitro flowering of *Spathiphyllum* in air-lift bioreactor cultures. Levels of electrical conductivity (EC), anions and cations, pH, ethylene, sugar content and photosynthetic enzymes were determined in bioreactor cultures of both flowering (FPs) and non-flowering (NFPs) plantlets over a growth period of 12 weeks. A decrease in ribulose 1,5-bisphosphate carboxylase/ oxygenase (Rubisco) activity with a corresponding increase in phosphoenolpyruvate carboxylase (PEPcase) activity occurred during floral induction of *Spathiphyllum* in vitro. Sucrose concentration decreased significantly in FPs, while no changes in glucose, fructose and total sugars were observed in both FPs and NFPs up to 8 weeks of culture. There were significant variations in mineral nutrient utilization between FP and NFP cultures. These results provide an insight to the physiological processes involved in inflorescence formation in *Spathiphyllum*

7

World Rabbit Sci.
(2008) 16: 157-164



DIETARY GRAPE POMACE AFFECTS LIPID PEROXIDATION AND ANTIOXIDATIVE STATUS IN RABBIT SEMEN

Eid Y.Z.

Department of Poultry Production, Faculty of Agriculture, Kafrelsheikh University,
33516 KaFr El-sHEiKH, Egypt.

Abstract

The objective of this study is to evaluate the effect of dietary grape pomace (GP) on certain characteristics, mainly lipid peroxidation and the antioxidative status of rabbit buck semen. Twenty seven adult New Zealand White rabbit bucks (6 months of age) were divided into three homogeneous groups (n=9) and randomly submitted to one of the three investigated dietary treatments. Animals in the first treatment (control) group were given the basal diet. The diets of the second (GP-10) and third (GP-20) treatment groups contained 10 and 20% of GP, respectively. Bucks received the experimental diets for 10 continuous weeks. GP did not appear to have any significant effect on body weight gain. Bucks receiving 10 or 20% dietary GP had a higher semen volume, 32% above that of rabbits in the control group ($P < 0.05$). The same trend was observed for sperm count. Dietary GP reduced the percentage of dead sperm and enhanced sperm motility. Interestingly, GP reduced lipid peroxidation in seminal plasma as indicated by TBARS, and significantly increased both the total antioxidant capacity and glutathione peroxidase activity ($P < 0.05$). In conclusion, the use of antioxidant dietary fibers rich in functional nutraceuticals (such as GP) may decrease lipid peroxidation and increase the antioxidative defense of rabbit semen.



8

International Journal of Horticultural Science
(2008) 14 (3): 39–44



Role of hydrogen peroxide and Pharmaplant-turbo against cucumber powdery mildew fungus under organic and inorganic production

Hafez, Y. M.1*, Bayoumi, Y. A.2¥, Pap, Z.2 & Kappel, N.2

1Dept. of Plant Pathophysiology, Plant Protection Institute, Hungarian Academy of Sciences, 1022, Herman Otto ut, 15, Budapest, Hungary 2Dept. of Vegetable and Mushroom Growing, Fac. of Hort. Corvinus University of Budapest, H-1118 Budapest, Villányi út 29–43, Hungary. *, ¥Permanent address: Faculty of Agriculture, Kafrelsheikh University,

Abstract

Cucumber leaves have been sprayed with a solution of hydrogen peroxide (H₂O₂) or Pharmaplant-turbo combined with organic or inorganic fertilizers under plastic house. Under the influence of H₂O₂, leaves exhibited resistance against *Podoshraera fusca* fungus, the causal agent of cucumber powdery mildew. H₂O₂ (15 mM) was able to decrease the disease severity from 90.4% to 12% in two experiments conducted in two seasons. Pharmaplant-turbo (Turbo) is a new chemical compound and used as an antifungal compound. Turbo in 1 ml/L was able to decrease the disease severity from 90.4% to 11.5% in both experiments as well. Both of H₂O₂ and Turbo were combined with organic treatment (compost + compost tea + seaweed extracts) which showed a significant effect against cucumber powdery mildew fungus and strongly suppressed it as compared to control leaves. Organic treatment produced higher vegetative growth characters and greater early and total yields as compared to inorganic treatment, also organic fruits produced the lower nitrate content and the higher ascorbic acid content as compared to inorganic fruits. Our study has indicated that, H₂O₂ and Turbo combined with organic fertilizers play a role in the resistance of cucumber against powdery mildew by decreasing the disease severity. We suggest to give more attention to the direct application of H₂O₂ in low concentration and Turbo against powdery mildew diseases and other plant diseases.

9

Journal of General Virology
(2008) 89: 799–808



Suppression of tobacco mosaic virus-induced hypersensitive-type necrotization in tobacco at high temperature is associated with down regulation of NADPH oxidase and superoxide and stimulation of dehydroascorbate reductase

L. Kira¹ly,¹³ Y. M. Hafez,²³ J. Fodor¹ and Z. Kira¹ly¹

¹Plant Protection Institute, Hungarian Academy of Sciences, PO Box 102, H-1525 Budapest, Hungary ²Department of Botany, Plant Pathology Branch, Faculty of Agriculture, Kafr-El-Sheikh University, Kafr-El-Sheikh, Egypt

Abstract

Tissue necroses and resistance during the hypersensitive response (HR) of tobacco to tobacco mosaic virus (TMV) are overcome at temperatures above 28 °C and the virus multiplies to high levels in the originally resistant N-gene expressing plants. We have demonstrated that chemical compounds that generate reactive oxygen species (ROS) or directly applied hydrogen peroxide (H₂O₂) are able to induce HR-type necroses in TMV-inoculated Xanthi-nc tobacco even at high temperatures (e.g. 30 °C). The amount of superoxide (O₂^{•-}) decreased, while H₂O₂ slightly increased in TMV- and mock-inoculated leaves at 30 °C, as compared with 20 °C. Activity of NADPH oxidase and mRNA levels of genes that encode NADPH oxidase and an alternative oxidase, respectively, were significantly lower, while activity of dehydroascorbate reductase was significantly higher at 30 °C, as compared with 20 °C. It was possible to reverse or suppress the chemically induced HR-type necrotization at 30 °C by the application of antioxidants, such as superoxide dismutase and catalase, demonstrating that the development of HR-type necroses indeed depends on a certain level of superoxide and other ROS. Importantly, high TMV levels at 30 °C were similar in infected plants, whether the HR-type necrotization developed or not. Suppression of virus multiplication in resistant, HR-producing tobacco at lower temperatures seems to be independent of the appearance of necroses but is associated with temperatures below 28 °C



Effectiveness of the antifungal black seed oil against powdery mildews of cucumber (*Podosphaera xanthii*) and barley (*Blumeria graminis* f.sp. *hordei*)


Yaser M. Hafez

Department of Botany (Plant Pathology Branch), Faculty of Agriculture, Kafrelsheikh University, Kafr-El-Sheikh, Egypt, and Plant Protection Institute, Hungarian Academy of Sciences, Budapest, Hungary

Abstract

When cucumber and barley leaves were sprayed with 0.5% black seed oil (BSO), rapeseed oil (R oil) and parafine oil (P oil), disease severity of the powdery mildew of cucumber (*Podosphaera xanthii*) was reduced from 52% (control) to 7.7% (BSO), 18.6% (R oil) and 20% (P oil). Similarly the disease severity of barley powdery mildew (*Blumeria graminis* f.sp. *hordei*) was greatly reduced from 63.4% (control) to 9.4% (BSO), 16% (R oil) and 16.4% (P oil). Oils inhibited the conidial germination of cucumber and barley powdery mildews to 29-30.7, 35-38 and 37-41% respectively, as compared to control (58-65%). Furthermore, mycelial growth of the pathogen was severely restricted after application of BSO and other oils. Levels of hydrogen peroxide (H₂O₂) and superoxide (O₂⁻) as well as the activity of the antioxidant enzymes in the treated leaves with oils and untreated (control) were measured and determined. H₂O₂ and O₂⁻ levels slightly increased, however some antioxidants are decreased such as dehydroascorbate reductase (DHAR) but other enzymes were increased such as ascorbate peroxidase (APX) and glutathione S transferase (GST). It can be concluded that the protective effect of oils against powdery mildews resulted mainly from the inhibition of conidial germination and suppression of the mycelial growth of the pathogens and there is slight activation of the host defence mechanisms. Therefore, it is important to giving more attention to BSO and other oils which have effectiveness against powdery mildew pathogens as an alternative control methods which safety and suitable for healthy and organic food production. Acta Biol Szeged 52(1):17-25 (2008)

11

Animal
(2008) 2 : 84–91

Ovarian follicular development, lipid peroxidation, antioxidative status and immune response in laying hens fed fish oil-supplemented diets to produce n-3-enriched eggs*

T. Ebeid¹, Y. Eid¹-, A. Saleh¹ and H. Abd El-Hamid²

¹Department of Poultry Production, Faculty of Agriculture, Kafrelsheikh University, 33516 Kafrelsheik, Egypt; ²Poultry Disease Department, Faculty of Veterinary Medicine, Alexandria University, Idfena, Egypt

Abstract

The objective of the present study was to research the effect of feeding laying hens fish oil-supplemented diets to produce n-3-enriched eggs on their ovarian follicular development, serum lipid peroxidation, antioxidative status and immune response. A total of 105 white Bovens hens at 24 weeks of age were housed in cages in an open-sided building under a 16 h light : 8 h dark lighting schedule. Birds were randomly divided into five treatments and were fed, ad libitum, diets containing 0%(control), 1.25%, 2.5%, 3.5% or 5.0% fish oil from 24 to 36 weeks of age. Egg production and weight were recorded. By weeks 35 and 36 of age 15 eggs were taken at random from each treatment to determine the yolk lipid profile and cholesterol content. At the end of the experimental period, 10 females from each treatment were randomly chosen, anaesthetised and killed by decapitation. Ovary and oviduct samples were immediately weighted and ovarian follicles were classified. Serum thiobarbituric acid-reactive substance (TBARS), hepatic TBARS and hepatic glutathione peroxidase (GSH-Px) activity were measured. No clear trend was observed concerning egg production and egg yolk cholesterol. As dietary fish oil levels increased, n-3-polyunsaturated fatty acids (n-3 PUFA) increased, whereas n-6 PUFA tended to decrease in yolk lipids. No negative effects were detected in ovary and oviduct weights, expressed in both absolute terms and relative to body weight. The numbers and total weights of large yellow follicles (LYF) in the ovary were not significantly affected by fish oil supplementation. Low levels(1.25% to 2.5%) of fish oil reduced both plasma and hepatic TBARS and enhanced GSH-Px activity. It is also interesting to note that inclusion of 2.5% fish oil in laying hen diets enhanced the antibody titre in laying hens. Therefore, it could be concluded that inclusion of fish oil in laying hen diets at moderate levels increased the n-3 fatty acids content in eggs, improved antioxidative status, enhanced the antibody response and did not have a negative influence on the different reproductive morphology parameters in laying hens.

12

International Journal of Horticultural Science
(2008) 14(3) :45 - 50

Using compost of grape manufacture and farm wastes as growing media in vegetable and ornamental nurseries

Bayoumi, Y. A. 1, El-Mahrouk, M. E.2, El-Aidy F.1 & Pap, Z.3

Horticulture Department, Faculty of Agriculture, Kafrelsheikh University of Egypt.
1Olericulture Branch 2 Floriculture Branch; 3 Dept. of Vegetable and Mushroom Growing,
Faculty of Horticulture, Corvinus University of Budapest, Hungary 1Current address:
Department. of Vegetable and Mushroom Growing, Faculty of Horticulture, Corvinus Univ.,
H-1118 Budapest, Villányi út 29–43., Hungary (Email: ybayoumi2002@yahoo.com.sg)

Abstract

This work was conducted at private nursery in Kafr El-Sheikh governorate during 2006 and 2007 seasons to investigate the possibility of using grape manufacture waste compost (GMWC) and farm wastes compost (FWC) in ornamental and vegetable nurseries as partially or totally replacement of coconut peat (CP) and vermiculite (V) in the growing medium and also to find out the optimum media of tomato (*Lycopersicon esculentum*, cv. Castle Rock) and Cockscomb (*Celosia plumosa*) as comparing to a mixture of CP and V (1:1 v/v). The authors used a ten mixtures as followed: 1- Control (CP+V at 1:1 v/v), 2- GMWC (100 %), 3- GMWC +CP (1:1 v/v), 4- GMWC + V(1:1 v/v),5- GMWC +CP + V (1:1:1 v/v/v), 6- FWC (100 %),7- FWC+ CP (1:1 v/v), 8- FWC+ V (1:1 v/v), 9- FWC+ CP+ V (1:1:1 v/v/v), 10- GMWC+FWC+CP+V (1:1:1:1 v/v/v/v). Data recorded as seedling height, No. of leaves, total pigments, shoot fresh and dry weights, root length and root fresh and dry weights in order to assess the quality of both transplants of tomato and cockscomb. Both seedlings grown in medium contain a mixture of GMWC+CP+V displayed quality traits similar or better as to those of recorded from the control treatment.

13

Acta Biologica Szegediensis
(2008) 52(1) :7-15



Improvement of postharvest keeping quality of white pepper fruits (*Capsicum annuum*, L.) by hydrogen peroxide treatment under storage conditions

Yousry A. Bayoumi

Horticulture Department, Faculty of Agriculture, Kafrelsheikh University, Kafr El-Sheikh, Egypt, and Department of Vegetable and Mushroom Growing, Faculty of Horticulture, Corvinus University, Budapest, Hungary

Abstract

Sweet pepper is one of the most important vegetable crops in the world, it has excellent nutritive value but it is susceptible to relatively fast quality changes after harvest time. The objective of the present research was to evaluate the effect of dipping pepper fruits in hydrogen peroxide solutions on postharvest keeping quality during storage time. Whole pepper fruits were soaked for 30 min in a solutions of hydrogen peroxide (0, 1, 5 and 15 mM) then, air dried and stored at room temperature (20°C) for 2 weeks and in fridge (10°C) for 4 weeks.

Hydrogen peroxide treatments significantly reduced weight loss, rot rate index and nitrate content of fruits specially with 15 mM hydrogen peroxide as compared with control treatment (0 mM hydrogen peroxide). Moreover, hydrogen peroxide treatments significantly increased general appearance, ascorbic acid content and the activity of the antioxidant enzymes such as ascorbate peroxidase and dehydroascorbate reductase. For dry matter and TSS%, there are no significant differences among treatments. Therefore, the use of hydrogen peroxide in postharvest treatments have a good potential strategy to improve the postharvest quality, extend shelf life period and maintained some nutritional quality as well as inhibited decay development of white peppers which natural infected under storage conditions. Acta Biol Szeged 52(1):7-15 (2008)



14

Crop Protection
(2008) 27 :653–659



Control of *Orobanche crenata* in legume intercropped with fenugreek (*Trigonella foenum-graecum*)

Mo´ nica Ferna´ ndez-Aparicioa, Amero A. Emeranb, Diego Rubialesa, _

aInstitute for Sustainable Agriculture, CSIC, Apdo. 4084, E-14080 Co´ rdoba, Spain

bDepartment of Agricultural Botany, Faculty of Agriculture, Kafr El-Sheikh University, 33516 Kafr El-Sheikh, Egypt

Abstract

Grain legume production in the Mediterranean Area is threatened by the holoparasitic plant *Orobanche crenata*, to which little resistance is available in affected crops. Control strategies have centred around agronomic practices and the use of herbicides, although success has been marginal. Several authors have described fenugreek as a suitable crop for intercropping with legumes, reducing the infection level of *O. crenata*; however, there is an important lack of experimental data and of a systematic research of the mechanisms involved in the reduction of parasitic infection. Two years of field experiments and further investigation by mini-rhizotron and pot experiments showed a decrease of *O. crenata* infection due to an allelopathic interference on the parasitic life cycle at the level of germination. Inhibition of *O. crenata* seed germination by allelochemicals released by fenugreek roots is suggested as the mechanism for reduction of *O. crenata* infection. r 2007 Elsevier Ltd. All rights reserved.

15

J. Phytopathology
(2008) 156 :419–424



Genetic Variation Among and Within *Uromyces* Species Infecting Legumes

A. A. Emeran¹, B. Román², J. C. Sillero², Z. Satovic³ and D. Rubiales⁴

Authors_ addresses: 1Faculty of Agriculture, Kafr El-Sheikh University, 33516 Kafr El-Sheikh, Egypt; 2IFAPA-CICE, Centro Alameda del Obispo, Apdo. 3092, 14080 Córdoba, Spain; 3Faculty of Agriculture, University of Zagreb, Svetosimunska 25, 10000 Zagreb, Croatia; 4Institute of Sustainable Agriculture, CSIC, Apdo. 4084, 14080 Córdoba, Spain (correspondence to D. Rubiales. E-mail: ge2ruozd@uco.es)

Abstract

Genetic variation of 30 different *Uromyces* isolates collected on faba bean, lentil, common vetch, pea, chickpea, alfalfa, cowpea and lupin was studied. Random Amplified Polymorphic DNA markers were used showing clear differences among *Uromyces* species. *Uromyces viciae-fabae* isolates clustered according to the host, with a clear cluster including all *U. viciaefabae* ex *Vicia faba* isolates. The *U. viciae-fabae* ex *Lens culinaris* isolate was the nearest to the cluster of *U. viciae faba* ex *V. faba* isolates, followed by *U. pisi* from Canada and *U. viciae-fabae* ex *V. sativa*. No association was found among molecular diversity and virulence or geographic origin within *U. vicia-fabae* ex *V. faba* isolates. Among the three *U. pisi* isolates considered, a great variability was observed and no grouping could be established. The most different isolate from the rest of species considered was *U. striatus*, followed by *U. vignae*. The two *U. ciceris-arietini* isolates clustered together and so did the two *U. lupinicolus* isolates.

16 Plant Pathology
(2007) 56 : 356



First report of sugar beet crown wart disease caused by *Urophlyctis leproides* in Egypt

M. I. Gouda^a and A. A. Emeran^b *

^aPlant Pathology Research, Institute Agriculture Research Center, Giza; and ^bPlant Pathology Branch, Department of Agricultural Botany, Faculty of Agriculture, University of Kafr El-Sheikh, 33516-Kafr El-Sheikh, Egypt

Abstract

Leaf and crown wart, marbled or beet root tumour are some of the common names given to a disease of sugar beet (*Beta vulgaris*) caused by *Urophlyctis leproides* (formerly *Physoderma leproides*). It was first reported in Algeria in 1894 (Trabut, 1894). Since then, the disease has been recorded in Argentina, Italy, Spain, United Kingdom, Palestine and the USA (Whitney, 1971). In Egypt, the disease was observed during harvest (2003, 2004, 2005) at three locations in the Nile Delta (Behira, Gharbia and Kafr El-Sheikh) on a range of sugar beet cultivars (Gloria, Kawmera, Oscar poly, Othus poly and Pleno). Disease incidence (DI) was less than 1% in 2003, 1–2% in 2004 and 3% in 2005. Although the DI is currently low, if the trend observed over the last few years continues, the disease could reach epidemic levels in the near future. The disease exhibited typical symptoms on root crowns and occasionally on petioles and leaf blades. Galls on leaf blades and petioles are greenish brown and have a rough appearance. Affected leaves are malformed. On the crown, galls range in size from 1 cm to (most commonly) 8–10 cm. These galls are spherical, rough and are attached to the host by a narrow base. Gall colour was variable, ranging from shades of green through yellow to brown depending on the age of plant cultivar. Galls occurred singly or coalesced to form complexes. Sections made (spores) surrounded by thickened wall. Sporangia were light brown, spherical to ovoid or concave ($25 \pm 5 \times 40 \pm 5 \mu\text{m}$) in diameter (Ruppel, 1995). As the gall decomposes, resting sporangia are released into the soil. Soil surveys were carried out in the affected areas and detected resting Twenty seedlings of *B. vulgaris* cv. Gloria were inoculated with 10^5 resting spores mL⁻¹ and incubated 22 ± 2 °C in continuous fluorescent light for 48 h, followed by 14 h light and 10 h dark. Uninoculated plants were used as a control. Symptoms appeared on the test plants after 11 days and sporangia were reisolated from the mature gall tissues (Mahmoody et al., 1997). No symptoms were observed on uninoculated controls. This is the first report of sugar beet crown wart disease (*U. leproides*) affecting *B. Vulgaris* in Egypt. sporangia.

17

Journal of Applied Horticulture
(2007) 9(2): 112-115



Rooting and growth response of grapevine nurslings to inoculation with arbuscular mycorrhizal fungi and irrigation intervals

Alaa El-din Khalil Omar

Department of Horticulture, Faculty of Agriculture, Kafr El-Sheikh University, Kafr El-Sheikh 33516, Egypt, E-mail: omaradks@yahoo.com

Abstract

This study was conducted during two successive seasons (2005 and 2006); in the experimental farm of Faculty of Agriculture, Kafr El Sheikh University; with the aim to investigate the influence of arbuscular mycorrhizal fungi (AMF) inoculation and irrigation intervals on growth of grapevine nurslings cv. Ruby King. Two mix mycorrhizal fungi including *Glomus fasciculatum* and *Glomus mosseae* were used for inoculation. The AMF inoculated and non-AMF nurslings were irrigated at 3, 6 and 9 days. The results showed that a combined treatment of AMF inoculation and irrigation at 3 days intervals recorded the highest values in terms of length of main root, total root length, root volume, root dry weight (%), top/root ratio, number of fine roots (< 2 mm), number of small roots (2-5 mm), number of leaves and leaf area per nursling. These results are of practical importance, as they highlight the potential of using mycorrhizal fungi inoculation for root development and growth improvement in grapevine nurslings and hence increases its adaptability upon transfer from the nursery to the open field.

18

Chemistry and Ecology
(2008) 24 :169–180

Biodegradation kinetics of cymoxanil in aquatic system

A.S. Derbalaha* and E.B. Belalb

aPesticides Department; bAgricultural Microbiology branch, Agricultural Botany Department, Faculty of Agriculture, Kafr El-Sheikh University, Kafr El-Sheikh, Egypt

Abstract

A potential method to detoxify pesticides in aquatic system is using bioremediation. In this study, four microorganisms (*Pseudomonas* sp (EB11), *Streptomyces* sp. (EB12), *Aspergillus niger* (EB13) and *Trichoderma viride* (EB14) were isolated from cucumber leaves previously treated with cymoxanil using enrichment technique. These strains were evaluated for their potential to detoxify cymoxanil in aquatic system at the concentration level of 5×10^{-4} M. The effect of pH and temperature on the growth ability of the tested strains was also investigated by measuring the intracellular protein and mycelia dry weight for bacterial and fungal strains, respectively. Moreover, the remaining toxicity of cymoxanil after 28 days of incubation with tested strains was evaluated to confirm the complete removal of any toxic materials (cymoxanil and its metabolites). The results showed that the optimum pH for the growth of cymoxanil degrading strains (bacteria and fungi) was 7. A temperature of 30°C appears to be the optimum for the growth of either fungal or bacterial strains. *Pseudomonas* sp. (EB11) was the most effective strain in cymoxanil degradation followed *Streptomyces* sp (EB12), *Trichoderma viride* (EB14) and *Aspergillus niger* (EB13), with half-lives of 4.33, 9.5, 17.3 and 24.7 days, respectively. The degradation of cymoxanil by bacterial strains was much faster than fungal one. There is no remaining toxicity of cymoxanil detected in aqueous media previously treated with *Pseudomonas* sp. (EB11) for 28 days. The results suggest that bioremediation by *Pseudomonas* sp. (EB11) are promising for the detoxification of cymoxanil in aqueous media.

19

hemosphere
(2008) 71 :2173–2179

Fate of imidacloprid in soil and plant after application to cotton seeds

Sherif E. El-Hamady a, R. Kubiak b, Aly S. Derbalah a,*

a Pesticides Department, Faculty of Agriculture, Kafr El-Sheikh University, 33516 Kafr-El-Sheikh, Egypt b Ecology Department, SLFA, Neustadt/W

Abstract

This study aimed to investigate the persistence of imidacloprid in soil after application to cotton seeds and to obtain a complete picture on the mass balance of this compound in soil and cotton plants. The study was carried out as a pot culture experiment under laboratory conditions using a Gaucho formulation containing ^{14}C -labeled imidacloprid. Three treatments of cotton seeds were made in sandy loamy soil: live seeds grown in autoclaved soil, dead seeds put in live soil and live seeds grown in live soil. Results showed that total ^{14}C recoveries decreased by time ranging 93.8–96.2, 77.1–88.4 and 53.5–62.4% of the applied radioactivity at 7, 14, and 21 d after application, respectively. The reduction in the extracted ^{14}C from soil coincided with the increase of non-extracted ones. Levels of bound ^{14}C was always less in autoclaved soil than in live ones. Results revealed also that only 1.8–6.8% of the applied ^{14}C was taken up by the plants and fluctuated within the test period. ^{14}C levels were higher in plants grown in autoclaved soil than those in live ones and the radioactivity tended to accumulate on the edges of cotton leaves. Most of the radioactivity in the soil extracts was identified as unchanged ^{14}C -imidacloprid. _ 2007 Elsevier Ltd. All rights reserved.

20

Journal of Environmental Sciences
(2008) 20 :262–267

Microbial detoxification of metalaxyl in aquatic system

Ahmed H. Massoud¹, Aly S. Derbalah^{1,*}, El-Sayed. B. Belal²

1. Pesticides Department, Faculty of Agriculture, Kafr El-Sheikh University. Kafr El-Sheikh 33516, Egypt. E-mail: masoudahm@yahoo.com

2. Agriculture Microbiology, Agriculture Botany Department, Faculty of Agriculture, Kafr El-Sheikh University, Kafr El-Sheikh 33516, Egypt

Abstract

Four microorganisms, *Pseudomonas* sp. (ER2), *Aspergillus niger* (ER6), *Cladosporium herbarum* (ER4) and *Penicillium* sp. (ER3), were isolated from cucumber leaves previously treated with metalaxyl using enrichment technique. These isolates were evaluated for detoxification of metalaxyl at the recommended dose level in aquatic system. The effect of pH and temperature on the growth ability of the tested isolates was also investigated by measuring the intracellular protein and mycelia dry weight for bacterial and fungal isolates, respectively. Moreover, the toxicity of metalaxyl after 28 d of treatment with the tested isolates was evaluated to confirm the complete removal of any toxic materials (metalaxyl and its metabolites). The results showed that the optimum degree pH for the growth of metalaxyl degrading isolates (bacterial and fungal isolates) was 7. The temperature 30° C appeared to be the optimum degree for the growth of either fungal or bacterial isolates. The results showed that *Pseudomonas* sp. (ER2) was the most effective isolate in metalaxyl degradation followed by *Aspergillus niger* (ER6), *Cladosporium herbarum* (ER4) and *Penicillium* sp. (ER3), respectively. There is no toxicity of metalaxyl detected in the supernatant after 28 d of treatment with *Pseudomonas* sp. (ER2). The results suggest that bioremediation by *Pseudomonas* sp. (ER2) isolate was considered to be effective method for detoxification of metalaxyl in aqueous media.

21

Land Contamination & Reclamation,
(2008) 16 (1) :13-23



Biodegradability of famoxadone by various microbial isolates in aquatic systems

A.S.H. Derbalah, E.B. Belal and A.H. Massoud

Department, Faculty of Agriculture, Kafr El-Sheikh Univ., Kafr El-Sheikh
33516, Egypt. Tel. 204-7325-5831, fax 204-7323-2032, email
aliderbalah@yahoo.com

Abstract

Bioremediation is defined as the process whereby organic wastes are biologically degraded under controlled conditions to an innocuous state, or to levels below concentration limits established by regulatory authorities. In this study, four microorganisms (*Pseudomonas* sp. (EB1), *Aspergillus niger* (EB2), *Cladosporium herbarum* (EB4) and *Penicillium* sp. (EB3) were isolated from cucumber leaves previously treated with famoxadone, using an enrichment technique. These isolates were evaluated for the detoxification of famoxadone at the concentration level of 100 $\mu\text{g mL}^{-1}$ in an aquatic system. The effects of pH and temperature on the growth of the tested isolates were also investigated by measuring the intracellular protein and mycelial dry weight for bacterial and fungal isolates, respectively. Moreover, the toxicity of famoxadone after 28 days of treatment with the tested isolates was evaluated to confirm the removal of any toxic materials (i.e. famoxadone and its metabolites). The results showed that the optimum pH for the growth of famoxadone-degrading isolates (i.e. bacterial and fungal isolates) was pH 7. A temperature of 30°C appears to be optimal for the growth of either fungal or bacterial isolates. Isolates of *Pseudomonas* sp. (EB1) and *Aspergillus niger* (EB2), were the most effective isolates in famoxadone degradation, followed by isolates of *Cladosporium herbarum* (EB4) and *Penicillium* sp. (EB3), respectively. A slight toxicity of famoxadone against *A. solani* as a sensitive target was detected in the supernatant treated by *Pseudomonas* sp. (EB1), *Aspergillus niger* (EB2) and *Penicillium* sp. (EB3) as compared with the control. The results suggest that bioremediation of famoxadone by *Pseudomonas* sp. (EB1), *Aspergillus niger* (EB2) and *Penicillium* sp. (EB3) is promising for the detoxification of famoxadone in aqueous media. Moreover, the tested microorganisms isolated from cucumber leaves may be able to significantly reduce the residue level of famoxadone in vegetable crops to below the maximum limit, especially under greenhouse conditions.

22

Contribuții Botanice
(2008) 43 : 135-140

MOLECULAR VARIABILITY OF EGYPTIAN AND HUNGARIAN BOTRYTIS CINEREA ISOLATES

EL-NAGGAR, Magdy^{1,2}, IRINYI Laszlo,²KÖVICS György Janos,
²SÁNDOR Erzsebet²

1 Kafr El-Sheikh University, Faculty of Agriculture, Agricultural Botany Department, Kafr El-Sheikh, Egypt

2 University of Debrecen, Centre of Agricultural Sciences, Department of Plant Protection, Debrecen, Hungary

e-mail: magdyelnagar77@yahoo.com

Abstract

An attempt was made to evaluate the variation between Hungarian and Egyptian isolates of *Botrytis cinerea* on the basis of molecular studies. Sequences from the minisatellites MSB1 and the widely used molecular marker in phylogenetic works, the translation elongation factor 1 subunit alpha (EF-1 α =tef1), were identified. The phylogenetic analysis was conducted with PAUP*4.0b by parsimony analysis. Topological robustness in parsimony analysis was estimated using 1000 bootstrap replicates. Our results show that two of the three studied Egyptian isolates of *B. cinerea* have been proved to be similar to the Hungarian ones on the basis of both minisatellites and tef1 sequences while the remaining one was completely different. The similarity between Hungarian and Egyptian isolates might be due to phylogenetic relationship or conidial dispersal between both countries.

23

Contribuții Botanice

(2008) 43 : 141-147



MYCOPARASITISM AND ANTAGONISTIC EFFICIENCY OF TRICHODERMA REESEI AGAINST BOTRYTIS SPP.

EL-NAGGAR Magdy^{1,2}, KÖVICS György Janos¹, SÁNDOR Erzsebet¹, IRINYI Laszlo¹

1. University of Debrecen, Centre of Agricultural Sciences, Department of Plant Protection, Debrecen, Hungary

2. Kafr El-Sheikh University, Faculty of Agriculture, Agricultural Botany Department, Kafr El-Sheikh, Egypt

e-mail: magdyelnaggar77@yahoo.com

Abstract

A preliminary study was made to evaluate the antagonistic efficiency and mycoparasitic activity of *Trichoderma reesei* against *Botrytis* species, viz. *B. cinerea*, *B. allii* and *B. fabae*. The bioassay method to evaluate the efficacy of *Trichoderma reesei* was the reduction of mycelial growth of *Botrytis* spp. in vitro. Significant effect was observed against *B. cinerea* and *B. fabae*, however, no effect was found against *B. allii*. Hyphal and sclerotial parasitism of *Trichoderma reesei* against *Botrytis* spp. were also studied. Coiling of hyperparasite fungus, penetration into *B. cinerea* hyphae, and stunt of the latest were observed as typical signs of mycoparasitism. Coiling of *B. allii* and stunted hyphae of *B. fabae* were the only signs of hyphal parasitism observed. The sclerotia of *B. cinerea* were also parasitized but sclerotia of *B. allii* and *B. fabae* were not



24

J Mater Sci
(2007) 42 :9717–9724



Elastic deformation of the rotating functionally graded annular disk with rigid casing

A. M. Zenkour

Department of Mathematics, Faculty of Education, Kafr
El-Sheikh University, Kafr El-Sheikh 33516, Egypt

Abstract

An accurate solution for a rotating functionally graded annular disk is presented. Material properties of the present annular disk are assumed to be graded in the radial direction according to a simple exponential-law distribution. The inner surface of the disk is pure metal whereas the outer surface of the disk is pure ceramic. The boundary condition of rigid casing is considered herein, that is the vanishing of the radial displacement at the outer surface. The boundary condition at the inner surface of the disk is taken to be vanishing either radial displacement or radial stress. Analytical solutions for the elastic deformation of the rotating functionally graded annular disks subjected to these boundary conditions are obtained. Numerical results for radial displacement, circumferential and radial stresses are presented. Comparisons between the different rotating homogeneous and functionally graded annular disks are made at the same angular velocity. The results show that distributions of stresses and displacement through the radial direction of the rotating annular disk vary with different parameters

25

J Mater Sci

(2008) 43 :2574–2589



Thermoelastic bending analysis of functionally graded sandwich plates

B. Ashraf M. Zenkour N. A. Alghamdi

Department of Mathematics, Faculty of Education,
Kafr El-Sheikh University, Kafr El-Sheikh 33516, Egypt

Abstract

The thermoelastic bending analysis of functionally graded ceramic–metal sandwich plates is studied. The governing equations of equilibrium are solved for a functionally graded sandwich plates under the effect of thermal loads. The sandwich plate faces are assumed to have isotropic, two-constituent material distribution through the thickness, and the modulus of elasticity, Poisson's ratio of the faces, and thermal expansion coefficients are assumed to vary according to a power law distribution in terms of the volume fractions of the constituents.

The core layer is still homogeneous and made of an isotropic ceramic material. Several kinds of sandwich plates are used taking into account the symmetry of the plate and the thickness of each layer. Field equations for functionally graded sandwich plates whose deformations are governed by either the shear deformation theories or the classical theory are derived. Displacement functions that identically satisfy boundary conditions are used to reduce the governing equations to a set of coupled ordinary differential equations with variable coefficients. The influences played by the transverse normal strain, shear deformation, thermal load, plate aspect ratio, side to thickness ratio, and volume fraction distribution are studied. Numerical results for deflections and stresses of functionally graded metal–ceramic plates are investigated.



26

Applied Mathematics & Information Sciences
(2008) 2(3) : 237–257



The Rotating Inhomogeneous Elastic Cylinders of Variable-Thickness and Density

M. N. M. Allam¹, A. M. Zenkour² and E. R. Elazab³

¹Department of Mathematics, Faculty of Science, Mansoura University
Mansoura 35516, Egypt E-mail address: mallam45@hotmail.com (M.N.M Allam)

²Department of Mathematics, Faculty of Education, Kafr El-Sheikh University
Kafr El-Sheikh 33516, Egypt E-mail address: zenkour@gmail.com (A.M. Zenkour)

³Department of Mathematics, Faculty of Science, Al-Azhar University
Assiut 71524, Egypt

Abstract

In this paper, an analytical solution is developed for the rotation problem of an inhomogeneous orthotropic cylinder with variable-thickness and density under plane strain assumption. The thickness of the cylinder and the elastic constants are taken as exponential functions in the radial direction but the density in a power law form. The cylinder may be solid or hollow with traction-free surface or clamped. On application of the boundary conditions, the stress and displacement for rotating homogeneous isotropic solid and hollow cylinders with uniform-thickness and density are obtained as special cases of the studied problem. Numerical results for stresses and displacement are presented in graphical forms. The effects of many parameters on stresses and displacement are investigated.

27

Appl. Math. Mech. -Engl. Ed
(2008) 29(12) :1601–1616

Elastic and viscoelastic solutions to rotating functionally graded hollow and solid cylinders

A. M. Zenkour^{1,2}, K. A. Elsibai³, D. S. Mashat¹

(1. Department of Mathematics, Faculty of Science, King Abdulaziz University, P. O. Box 80203, Jeddah 21589, Saudi Arabia;

2. Department of Mathematics, Faculty of Education, Kafr El-Sheikh University, Kafr El-Sheikh 33516, Egypt;

3. Department of Mathematics, Faculty of Science, Mansoura University, Mansoura 35516, Egypt)
(Communicated by GUO Xing-ming)

Abstract

Analytical solutions to rotating functionally graded hollow and solid long cylinders are developed. Young's modulus and material density of the cylinder are assumed to vary exponentially in the radial direction, and Poisson's ratio is assumed to be constant. A unified governing equation is derived from the equilibrium equations, compatibility equation, deformation theory of elasticity and the stress-strain relationship. The governing second-order differential equation is solved in terms of a hypergeometric function for the elastic deformation of rotating functionally graded cylinders. Dependence of stresses in the cylinder on the inhomogeneous parameters, geometry and boundary conditions is examined and discussed. The proposed solution is validated by comparing the results for rotating functionally graded hollow and solid cylinders with the results for rotating homogeneous isotropic cylinders. In addition, a viscoelastic solution to the rotating viscoelastic cylinder is presented, and dependence of stresses in hollow and solid cylinders on the time parameter is examined.



28

Hindawi Publishing Corporation
International Journal of Photoenergy
2008 Volume: 1-8



Two-Dimensional Free Energy Surfaces for Electron Transfer Reactions in Solution

Shigeo Murata,¹ Maged El-Kemary,² and M. Tachiya¹

¹National Institute of Advanced Industrial Science and Technology (AIST), AIST Tsukuba Central 5, 1-1-1 Higashi, Tsukuba 305-8565, Japan

²Chemistry Department, Faculty of Education, Kafra ElSheikh University, Kafra ElSheikh 33516, Egypt
Correspondence should be addressed to Shigeo Murata, shigeo.murata@aist.go.jp

Abstract

Change in intermolecular distance between electron donor (D) and acceptor (A) can induce intermolecular electron transfer (ET) even in nonpolar solvent, where solvent orientational polarization is absent. This was shown by making simple calculations of the energies of the initial and final states of ET. In the case of polar solvent, the free energies are functions of both D-A distance and solvent orientational polarization. On the basis of 2-dimensional free energy surfaces, the relation of Marcus ET and exciplex formation is discussed. The transient effect in fluorescence quenching was measured for several D-A pairs in a nonpolar solvent. The results were analyzed by assuming a distance dependence of the ET rate that is consistent with the above model. Copyright © 2008 Shigeo Murata et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

29

LMSA_A_345542 803.cls

(2008) 25 1:25



Immobilization of Urease on (HEMA/IA) Hydrogel Prepared by Gamma Radiation

SOHA M. HAMDY¹, SAMIA EL-SIGENY² and MANAL F. ABOU TALEB³

¹Biochemistry Division, Faculty of Science, Fayoum University, Egypt

²Chemistry and Physics Department, Faculty of Education, Kafr El-Sheikh University, Egypt

³National Center for Radiation Research and Technology, P.O. Box 29, Nasr City, Cairo, Egypt

Abstract

In the present study, the copolymeric hydrogels based on 2-hydroxyethyl methacrylate (HEMA) and itaconic acid (IA) were synthesized by gamma radiation induced radical polymerization, in order to examine the potential use of these hydrogels in immobilization

of *Citrullus vulgaris* urease. Gelation and Swelling properties of PHEMA and copolymeric P (HEMA/IA) hydrogels with different IA contents (96.5/3.5, 94.4/5.6 and 92.5/7.5 mol) were studied in a wide pH range. Initial studies of so-prepared hydrogels show interesting pH sensitivity in swelling and immobilization. *C. vulgaris* urease was immobilized on HEMA/IA (92.5/7.5) at 6 kGy with 41.3% retention of activity. The properties of free and immobilized urease were compared. Immobilized urease maintained a higher relative activity than free urease at both lower and higher pH levels, indicating that the immobilized urease was less sensitive to pH changes than the free urease. The K_m value of the immobilized urease was approximately 2 times higher than that of the free urease. Temperature stability was improved for immobilized enzyme. The free form exhibited a loss about 80% of activity upon incubation for 15 min at 80°C. The influence of various heavy metal ions at the concentration of 1 mM was improved after enzyme immobilization. The immobilization of *C. vulgaris* urease on HEMA/IA (92.5/7.5) at 6 kGy showed a residual activity of 47 % after 4 reuses.

3. LMSA_A_351712 803.cls
(2008) 46 : 1-9



Hybrid Nanocomposite Prepared by Graft Copolymerization of 4-Acryloyl morpholine onto Chitosan in the Presenc of Organophilic Montmorillonite

SAMIA AL-SIGENY,¹ MANAL F. ABOU TALEB^{2,*} and NABIL A. EL-KELESH²

¹Chemistry and Physics Department, Faculty of Education at Kafr El-Sheikh, Kafr El-Sheikh University, Egypt

²National Center for Radiation Research and Technology, P.O. Box 29, Nasr City, Cairo, Egypt

Abstract

Organophilic montmorillonite (OMMT) was synthesized by cationic exchange between Na-MMT and Vinyl benzyl triphenyl phosphonium chloride in an aqueous solution. A new nanocomposite consisting of 4-acryloyl morpholine-chitosan and OMMT was prepared by γ -ray irradiation polymerization. The intercalation spacing of these nanocomposites was investigated with X-ray diffraction and its thermal stabilities by adding nanocomposites were characterized by thermal gravimetric analysis. The nanocomposites showed improved resistance to water absorption. The most interesting application of the nanocomposite is its ability for adsorption purification of waste water containing acid dyes. One of the objectives in this study was to develop new and active prepared copolymers which can be examined for their antimicrobial activities. It was found that the copolymer nanocoposite based on phosphonium group and some heavy metal ions in its structure having broad spectrum against pathogenic bacteria such as *Staphylococcus aureus*, *Escherichia coli* and *Aspergillus flavus* fungi

31 Reactive & Functional Polymers
(2008) 68 : 1680–1693



Convenient approaches for the synthesis and characterization of well-defined linear-dendritic diblock copolymers having a definite number of peripheral primary amino groups: Exact control of numbers introduced and dendritic distribution to enhance co-operative effect

Ashraf A. El-Shehawy a,* , Kenji Sugiyama b, Akira Hirao b

a Chemistry Department, Faculty of Education, Kafr El-Sheikh University, Kafr El-Sheikh 33516, Egypt

b Organic and Polymeric Materials Department, Graduate School of Science and Engineering, Tokyo Institute of Technology, 2-12-1 Ohokayama, Meguro-ku, Tokyo 152-8552, Japan

Abstract

Efficient and convenient approaches have been developed to prepare a wide array of well-defined linear-dendritic diblock copolymers having a definite number of peripheral primary amino groups by utilizing linear polystyrenes end-functionalized with a definite number of dendritic benzyl bromide and/or benzyl alcohol functionalities as precursory polymers. Various approaches including modified Gabriel synthesis, Michael addition and alkylation reactions have been developed for their synthesis. The resulting primary amine-end-functionalized copolymers always exhibited SEC profiles showing asymmetric long-tailed distributions and their true molar masses were obtained after the acetylation of their peripheral dendritic amino groups with acetyl chloride. All the resulting copolymers possessed well-defined linear-dendritic diblock architectures, with controlled molecular weights. In these polymer series, the molecular weight of linear polystyrene chains remained almost unchanged ($M_n \approx 19.3$ kg/mol), while the dendritic blocks increased in both size and molecular weight as the number of peripheral dendritic end-functional groups was increased. The resulting copolymers were fully characterized by elemental, FT-IR, TLC-FID, GPC, 1H and ^{13}C NMR analyse



32

Tetrahedron: Asymmetry
(2008)19 : 425–434



Application of well-defined chain-end functionalized polystyrenes with dendritic chiral ephedrine moieties as reagents for highly catalytic enantioselective addition of dialkylzincs to aldehydes

Ashraf A. El-Shehawy,* Kenji Sugiyama and Akira Hirao

Department of Chemistry, Faculty of Education, Kafr El-Sheikh University,
Kafr El-Sheikh, PO Box 33516, Egypt.

Organic and Polymeric Materials Department, Graduate School of Science and Engineering, Tokyo Institute of Technology,
2-12-1, Ohokayama, Meguro-ku, Tokyo 152-8552, Japan

Abstract

A series of well-defined chain-end-functionalized polystyrenes having a definite number of chiral ephedrine moieties dendritically distributed at the periphery of their hyperbranched chain-ends were evaluated as chiral catalysts for the enantioselective addition of dialkylzinc reagents to aldehydes. These dendritic macromolecules worked well as homogeneous chiral catalysts and exhibited high catalytic activity and enantioselectivity very similar to those observed for the corresponding monomeric chiral catalysts. The optimum amount of chiral catalyst was found to be 5 mol %. A profound number effect of the chiral ephedrine moieties was observed, and PS(Ephed)8 having eight chiral ephedrine moieties at the periphery was found to be superior to other dendritic chiral catalysts. The enantioselectivity reached a value of 95% in the addition of diisopropylzinc to 3-phenylpropanal. The dendritic chiral catalysts could be easily recovered from the reaction solution by using a solvent precipitation method, and the recovered catalyst showed no significant loss of its catalytic activity or enantioselectivity. _ 2008 Elsevier Ltd. All rights reserved.

33

hem. Pharm. Bull.
(2008) 56 (12): 1651-1657

Development and Validation of an Adsorptive Stripping Voltammetric Method for the Quantification of Vincamine in Its Formulations and Human Serum Using a Nujol-Based Carbon Paste Electrode

Amr Mohamed Beltagy

Chemistry & Physics Department, Faculty of Education, Kafr El-Sheikh University; 33516 Kafr El-Sheikh, Egypt.

Abstract

An easy, rapid and selective adsorptive stripping voltammetry (AdSV) method for the determination of vincamine in its formulation and human serum was developed and validated. It was based on the oxidation of the drug onto a Nujol-based carbon paste electrode. The stripping step was carried out by using a square-wave (SW) potential-time voltammetric excitation signal. The optimal experimental variables as well as accumulation parameters were investigated as; frequency f_{120} Hz, scan increment DEi_{10mV} , pulse-amplitude DEa_{25mV} and an accumulation potential E_{acc} of 0.0 V using a Britton–Robinson (B-R) universal buffer of pH 5 as a supporting electrolyte. After validation of the described method, it was applied for determination of vincamine in its formulation and human serum. Mean recovery of $100.41_{0.74}$ (n_5) was achieved for assay of vincamine in Oxybral® capsules. Limits of detection and quantitation of 6.0_{10_9M} (2.20 ng ml $_{-1}$) and 2_{10_8M} (7.33 ng ml $_{-1}$) vincamine were achieved in human serum with a mean recovery of $99.5_{1.79}\%$, without prior extraction of the drug. No interferences were observed in formulation and/or human serum. Due to high sensitivity and specificity of the developed method, it was successfully applied for evaluating some pharmacokinetic parameters of two healthy volunteers after administration of a single oral Oxybral® capsule.

34

J Appl Electrochem
Published on line

Simultaneous determination of trace aluminum (III), copper (II) and cadmium (II) in water samples by square-wave adsorptive cathodic stripping voltammetry in the presence of oxine

A. M. Beltagi M. M. Ghoneim

Chemistry & Physics Department, Faculty of Education,
Kafr El-Sheikh University, 33616 Kafr El-Sheikh, Egypt
Received: 20 July 2008 / Accepted: 21 October 2008
_ Springer Science+Business Media B.V. 2008

Abstract

A validated adsorptive cathodic stripping voltammetry method is described for simultaneous determination of Al(III), Cu(II) and Cd(II) in water samples. In acetate buffer (pH 5) containing 10 IM oxine, these metal ions were determined as oxine complexes following adsorptive accumulation onto the HMDE at -0.05 V versus Ag/AgCl/KCl. The best signal to noise ratio was obtained using a square wave of scan increment 10 mV, frequency 120 Hz, and pulse-amplitude 25 mV. Limits of detection as low as 0.020 $\mu\text{g L}^{-1}$ Al(III), 0.012 $\mu\text{g L}^{-1}$ Cu(II) and 0.028 $\mu\text{g L}^{-1}$ Cd(II) were achieved. Interference due to various cations (K(I), Na(I), Mg(II), Ca(II), Mn(II), Fe(III), Bi(III), Sb(III), Se(IV), Pb(II), Zn(II), Ni(II), Co(II)), anions (Cl⁻, NO₃⁻, SO₄²⁻, PO₄³⁻) and ascorbic acid was minimal as the measured signals change by 4% at the maximum. The stripping voltammetry method was successfully applied for simultaneous determination of Al(III), Cu(II) and Cd(II) in tap and natural bottled water samples.

35

Talanta

(2008) 74 :851–859



Development of a voltammetric procedure for assay of the antihistamine drug hydroxyzine at a glassy carbon electrode: Quantification and pharmacokinetic studies

A.M. Beltagi a,* , O.M. Abdallah b, M.M. Ghoneimc

a Chemistry & Physics Department, Faculty of Education, Kafr El-Sheikh, University , 33516 Kafr El-Sheikh, Egypt

b Analytical Chemistry Department, Faculty of Pharmacy (Girls), Al-Azhar University, Nasr City, Cairo, Egypt

c Analytical Chemistry Research Unit, Chemistry Department, Faculty of Science, Tanta University, 31527 Tanta, Egypt

Abstract

An electrochemical study of hydroxyzine at a glassy carbon electrode was carried out in the Britton–Robinson universal buffer of pH 2–11. Hydroxyzine was oxidized in a single two-electron irreversible process controlled mainly by adsorption. A simple, sensitive and time-saving squarewave adsorptive anodic stripping voltammetric procedure has been developed for determination of hydroxyzine in its commercial tablets and human serum without prior extraction. The optimized procedural conditions were: frequency = 120 Hz, scan increment = 10mV, pulse-amplitude = 25mV, accumulation potential = -0.3V, accumulation time = 90–300 s and a Britton–Robinson universal buffer of pH 4 as a supporting electrolyte. Mean recoveries of 100.5 ± 0.71 and $98.6 \pm 1.12\%$ ($n = 5$) were achieved for assay of hydroxyzine in Atarax® 10 and 25 mg dosage forms, respectively. Limit of detection of 1.5×10^{-8} mol L⁻¹ (5.624 ngmL⁻¹) and limit of quantitation of 5.0×10^{-8} mol L⁻¹ (18.746 ngmL⁻¹) were achieved in human serum with a mean recovery of $98.4 \pm 1.22\%$, without prior extraction of the drug. Moreover, the described procedure was applied for evaluating the pharmacokinetic parameters of hydroxyzine in plasma of two healthy volunteers after administration of a single oral dose (Atarax®—25 mg). © 2007 Elsevier B.V. All rights reserved.

36

Micropaleontology
(2007) 53 : 127-144

Nannofossil biostratigraphy of the Paleocene-lower Eocene succession in the Thamad area, east central Sinai, Egypt

Mahmoud Faris¹ and Aziz Mahmoud Abu Shama²

¹Geology Department, Faculty of Science, Tanta University, Egypt
email: mhmfaris@yahoo.com

²Biological and Geological Sciences Department, Faculty of Education Kafr El Sheikh University, Egypt
,email: mabushama2002@yahoo.com

Abstract

In the Thamad area, east central Sinai, two stratigraphic sections are measured and sampled; Gebel El Mishiti and G. El Keeh, in order to determine the nannofossil biostratigraphy of the Paleocene-lower Eocene rocks. Gebel El Mishiti section is considered one of the most complete sections across the K/P boundary in Egypt where all the exposed rocks at G. El Keeh belong to the lower

Eocene Egma Formation. A nearly complete succession of Paleocene nannofossil biozones is recorded from the Esna Formation at G. ElMishiti. They are arranged from base to top as follows: *Markalius inversus* (NP1), *Cruciplacolithus tenuis* (NP2), *Chiasmolithus danicus* (NP3), *Ellipsolithus macellus* (NP4), *Fasciculithus tympaniformis* (NP5), *Heliolithus kleinpellii* (NP6), *Discoaster mohleri* (NP7/8) and *Discoaster multiradiatus* (NP9) zones. The Danian/Selandian boundary is located at the base of Zone NP5 at the level of the first appearance of *Fasciculithus* taxa. The Selandian/Thanetian boundary can be traced tentatively at the base of the *Discoaster mohleri* Zone (NP7/8). The Paleocene/Eocene boundary is traced between the NP9a/NP9b subzonal boundary which is marked by the first appearances of *Discoaster araneus*, *Rhomboaster calcitraba*, *R. bitrifida* and *R.cuspis*. The Paleocene /Eocene boundary lies within the upper part of the Esna Formation. A very thin layer of conglomeratic chalk with no paleontological break is located between Zone NP9 and NP10. However, a very short nannofossil break is noted within Zone NP10 (subzones NP10b and NP10c are missing) which is overlain by Zone NP11. The

lower part of the latter covers the topmost part of the Esna Formation where its upper part lies within the base of the Egma Formation; therefore, a conformable relationship between the two formations is suggested. At Gebel El Mishiti, the lower Eocene nannofossil biozones NP11, NP12, NP13 and lower/middle Eocene Zone NP14 are recorded. On the other hand, Gebel El Keeh section, which is about 95m thick is only represented by the *Discoaster subloadoensis* Zone (NP14).

37

Parasitology Research

Founded as Zeitschrift für Parasitenkunde

(2009) 104(4) : 745-752



The first detection of *Cryptosporidium* deer-like genotype in cattle in Japan

Said Amer^{1, 2}, Hajime Honma¹, Makoto Ikarashi¹, Ryu Oishi¹, Mikiko Endo¹, Kenichi Otawa¹ and Yutaka Nakai¹

(1) Laboratory of Sustainable Environmental Biology, Graduate School of Agricultural Science, Tohoku University, 232-3 Yomogita, Naruko-onshen, Osaki Miyagi, 989-6711, Japan

(2) Present address: Department of Biological and Geological Sciences, Faculty of Education, Kafar El Sheikh University, Kafar el-Sheikh, 33516, Egypt

Abstract

The general perception is that cattle are major reservoirs for *Cryptosporidium parvum* infections in humans and that *C. parvum* is a major cause of diarrhea and production loss in cattle. Adult cattle may play an important role as cryptic carrier of the infection. *Cryptosporidium* spp. in asymptomatic adult dairy cattle from some farms around Osaki area, Miyagi prefecture, Japan, was examined on a field visit during August, 2007, by polymerase chain reaction techniques for detection, genotyping, and subtyping. *Cryptosporidium* oocysts were detected in the feces of five out of 50 animals. Of the five *Cryptosporidium*-positive specimens available for molecular analysis, *C. parvum* was identified in three specimens, *Cryptosporidium* deer-like genotype in one, and *Cryptosporidium andersoni* in one specimen. Amplification of *Cpgp60* from *C. andersoni* and *Cryptosporidium* deer-like genotype samples revealed that these samples have light concurrent *C. parvum* infection. Sequence analysis of the 60-kDa glycoprotein gene indicated that all *C. parvum* samples are IIa subtype. Detection of *Cryptosporidium* deer-like genotype is geographically unique in Japan. The genetic diversity of *Cryptosporidium* in dairy cattle in Japan may be much greater than that reported before.



Downregulation of male-specific cytochrome P450 by profenofos

Gihan G. Moustafa^{1, 2, a)}, Zein S. Ibrahim^{1, 3, a)}, Mohamed M. Ahmed⁴⁾, Mervat H. Ghoneim²⁾, Kentaro Q. Sakamoto¹⁾, Mayumi Ishizuka^{1,*}) and Shoichi Fujita¹⁾

1)Laboratory of Toxicology, Department of Environmental Veterinary Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

2)Department of Forensic Medicine and Toxicology, Faculty of Veterinary Medicine, Zagazig University, Zagazig 44519, Egypt.

3)Department of Biochemistry and Physiology, Faculty of Veterinary Medicine, Kafr El-Sheikh University, Kafr El-Sheikh 54363, Egypt

4)Laboratory of Biochemistry, Department of Biomedical Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

Abstract

The health hazards of individual organophosphorus insecticides have been characterized by their acute toxicity, mainly by investigating their cholinesterase inhibition. However, the chronic effects of most of these toxicants on the drug-metabolizing enzymes have not been investigated. Profenofos (O-4-bromo-2-chlorophenyl O-ethyl S-propyl phosphorothioate) is an organophosphorus pesticide widely used in cotton cultivation. In the present study, we investigated the effect of profenofos on male-specific cytochrome P450 (CYP) enzymes in adult Wistar rats. We orally administered 17.8 mg/kg body weight, twice weekly for 65 days. Profenofos downregulated levels of hepatic and testicular CYP2C11 and CYP3A2 mRNA and protein expression. Testicular aromatase (CYP19A) mRNA was decreased in the profenofos-treated rats compared to controls. Overall, the present study suggests that profenofos acts as an endocrine disruptor of male-specific CYP enzymes and affects testosterone concentration, which implicates its deleterious effects on animal or human males chronically exposed to organophosphorus pesticide.

39

J BIOCHEM MOLECULAR TOXICOLOGY

(2008) 22 : 77-84



The Induction of Cytochrome P450 1A1 by Sudan Dyes

Nahla A. G. Ahmed Refat,^{1,2} Zein Shaban Ibrahim,^{1,3} Gihan Gamal Moustafa,^{1,4}
Kentaro Q. Sakamoto,¹ Mayumi Ishizuka,¹ and Shoichi Fujita¹

Department of Pathology, Faculty of Veterinary Medicine, Zagazig University, Egypt³Department of Biochemistry and Physiology, Faculty of Veterinary Medicine, Kafre El-Sheikh University, Egypt
Department of Forensic Medicine and Toxicology, Faculty of Veterinary Medicine, Zagazig University, Egypt

Abstract

Azo dyes form a major class of chemically related compounds that are ubiquitous in foods, paints, printing inks, cosmetics, and also used as biological stains in histological and histopathological laboratories and clinics. Sudan I, sudan III, and sudan IV have been classified as category 3 carcinogens by International Agency for Research on Cancer. In this study, we investigated the difference between these three sudan dyes in induction of CYP1A1. We intraperitoneally treated Wistar rats with each of the three sudan dyes (I, III, and IV) for 3 days. Treatment of Wistar rats with sudan I produced the highest induction of CYP1A1 protein and mRNA whereas treatment of Wistar rats with sudan III produced about two third of CYP1A1 protein and mRNA than induced by sudan I. Furthermore, treatment of Wistar rats with sudan IV produced the lowest induction of CYP1A1 protein and mRNA which is about two third of that induced with sudan III treatment. We further investigated the effect of these sudan dyes on CYP1A1 transcription through investigating the xenobiotic response element (XRE) reporter activity in HepG2. The XRE reporter activity study showed the same trend of activity of sudan dyes comparable to the effects on CYP1A1 mRNA and protein. Immunohistochemical study revealed a differential pattern of distribution of CYP1A1 protein in rat liver among the three sudan dyes, apparent in the centrilobular and midzonal region with sudan III, progressing to panlobular with sudan I, whereas sudan IV showed a reversal of pattern of induction with the most intense staining in the periportal region. Our results suggest that there is an inverse relationship between the molecular size of the three sudan dyes and their ability to induce CYP1A1.

40

APPLIED AND ENVIRONMENTAL
MICROBIOLOGY
(2007) 73 : 6686–6690



Zoo Animals as Reservoirs of Gram-Negative Bacteria Harboring Integrons and Antimicrobial Resistance Genes

Ashraf M. Ahmed,^{1,3} Yusuke Motoi,¹ Maiko Sato,¹ Akito Maruyama,¹
Hitoshi Watanabe,²
Yukio Fukumoto,² and Tadashi Shimamoto^{1*}

Laboratory of Food Microbiology and Hygiene, Graduate School of Biosphere Science, Hiroshima University, Higashi-Hiroshima 739-8528, Japan¹; Hiroshima City Asa Zoological Park, Asa-cho Asakita-ku, Hiroshima 731-3355, Japan²; and Department of Microbiology, Faculty of Veterinary Medicine, Kafr El-Sheikh University, Kafr El-Sheikh 33516, Egypt³

Abstract

A total of 232 isolates of gram-negative bacteria were recovered from mammals, reptiles, and birds housed at Asa Zoological Park, Hiroshima prefecture, Japan. Forty-nine isolates (21.1%) showed multidrug resistance phenotypes and harbored at least one antimicrobial resistance gene. PCR and DNA sequencing identified class 1 and class 2 integrons and many β -lactamase-encoding genes, in addition to a novel AmpC β -lactamase gene, blaCMY-26. Furthermore, the plasmid-mediated quinolone resistance genes qnr and aac(6_{1b})-Ib-cr were also identified.

41

Journal of Antimicrobial Chemotherapy
(2007) 59 :184–190



Proteus mirabilis clinical isolate harbouring a new variant of Salmonella genomic island 1 containing the multiple antibiotic resistance region

Ashraf M. Ahmed^{1,2}, Amjad I. A. Hussein³
and Tadashi Shimamoto^{1*}

¹Laboratory of Food Microbiology and Hygiene, Graduate School of Biosphere Science, Hiroshima University,

Higashi-Hiroshima 739-8528, Japan; ²Department of Microbiology, Faculty of Veterinary Medicine, Kafr El-Sheikh University, Kafr El-Sheikh 33516, Egypt; ³Graduate School of Medical Science, Kanazawa University, Kanazawa 920-8640, Japan

Received 4 September 2006; returned 12 October 2006; revised 23 October 2006; accepted 25 October 2006

Abstract

Objectives: A clinical isolate of *Proteus mirabilis* strain 18306, which displayed the multidrug resistance phenotype of *Salmonella* genomic island 1 (SGI1), was examined for the presence of this island including its multiple antibiotic resistance genomic region. **Methods:** *P. mirabilis* 18306 was isolated in March 2006 from a patient in Palestine with diabetic foot infection. Antibiotic susceptibility tests and various molecular techniques, including PCR, cloning and DNA sequencing were used for detection and characterization of SGI1 in *P. mirabilis* 18306. **Results:** *P. mirabilis* 18306 showed the typical multidrug resistance phenotype of SGI1 as it was resistant to ampicillin, chloramphenicol, streptomycin, sulphonamides and tetracycline, in addition to trimethoprim and nalidixic acid. Molecular characterization showed that *P. mirabilis* 18306 harboured a structure similar to SGI1, except that the *aadA2* gene, which confers resistance to streptomycin and spectinomycin, of standard SGI1 had been replaced with *dfrA15*, which confers resistance to trimethoprim. Furthermore, the nucleotide sequence of the extrachromosomal circular form of SGI1 in *P. mirabilis* was found to be identical to that of *Salmonella* Typhimurium DT104. However, PCR results showed that *P. mirabilis* 18306 was negative for the left and right junctions which represent the integration sites of SGI1 into *Salmonella enterica* chromosome. Hence, this new variant of SGI1 may be integrated at a different site into the chromosome of *P. mirabilis* 18306. Tn1826-derived class 2 integron, which carries only two gene cassettes, *sat2* and *aadA1*, was also identified in this strain. **Conclusions:** In this study, we identified a new variant SGI1 containing the multiple resistance genomic region in a multidrug-resistant strain of *P. mirabilis*. This is the first report for SGI1 in a genus other than *Salmonella*.



42

International Journal of Antimicrobial Agents
(2008) 32 :256–261



Emergence of a cefepime- and ceftazidime-resistant *Citrobacter freundii* clinical isolate harbouring a novel chromosomally encoded AmpC β -lactamase, CMY-37 Ashraf M. Ahmed¹, Tadashi Shimamoto*

Department of Bacteriology, Faculty of Veterinary
Medicine, Kafr El-Sheikh University, Kafr El-Sheikh 33516, Egypt.
Laboratory of Food Microbiology and Hygiene, Graduate School of Biosphere Science, Hiroshima
University, Higashi-Hiroshima, Japan

Abstract

Citrobacter freundii strain 4306 was isolated from a urine specimen of a patient in March 2006 in Palestine. This strain showed a unique multidrug resistance phenotype, as it was resistant both to 7- β -methoxy- and oxyimino-cephalosporins, including cefepime, ceftazidime and monobactams, in addition to quinolones, streptomycin and trimethoprim/sulfamethoxazole. Clavulanic acid did not act synergistically with cephalosporins by the double-disk synergy test. Molecular characterisation showed that the resistance to 7- β -methoxy and oxyimino-cephalosporins was due to a novel AmpC β -lactamase, designated CMY-37, with an isoelectric point of \sim 9.0. CMY-37 is a variant of *C. freundii* chromosomal AmpC enzymes with at least seven amino acid substitutions. One of these substitutions, L316I, is located within the R2 loop that is considered the hotspot region responsible for the extended substrate spectrum in class C β -lactamases. The blaCMY-37 gene was cloned and expressed in *Escherichia coli* TG1. CMY-37 is chromosomally encoded and is not associated with ISEcp1-like element. Phylogenetic analysis suggested that CMY-37 is the origin of many plasmid-mediated AmpC β -lactamases. This study highlights the emergence of cefepime and ceftazidime resistance in *C. freundii* owing to a new type of AmpC β -lactamase



43

Trop Anim Health Prod
Published on line



Ocular infection of donkeys (*Equus asinus*) with *Setaria equine*

Mohamed A. Marzok & Abdel-Razek Y. Desouky

Department of Surgery, Faculty of Veterinary Medicine,
Kafr El-Sheikh University,
Kafr El-Sheikh 33516, Egypt

Abstract

Abstract Seven donkeys raised in different locations in Egypt were found to have a unilateral eye showing motile white worms in the aqueous humor. The parasites were surgically removed from the anterior chamber of the eye in five out of the seven donkeys, and were cleared, mounted and identified as *Setaria equina* based on light microscopic features. The ocular infection with *S. equina* reported herein may be the first reported aberrant cases in the Egyptian animals.

44 Trop Anim Health Prod
(2008) 40 :419–426



Cryptosporidiosis in buffalo calves (Bubalus bubalis): Prevalence and potential risk factors

Sabry A. El-Khodery & Salama A. Osman

Department of Animal Medicine, Faculty of Veterinary
Medicine, Kafri El-Sheikh University,
Kafri El-Sheikh, Egypt

Abstract

The objective of the present study was to describe the prevalence and risk factors associated with cryptosporidiosis in buffalo calves in Middle Egypt. During one year, 458 fecal samples were collected from buffalo calves less than 3 month age in 55 small scale herds and examined for the presence of *Cryptosporidium* oocysts. Data describing age, gender, season, and herd management practices were gathered to assess potential risk factors. Fecal examination showed that 14.19% of the examined calves were positive for *Cryptosporidium* spp. Calves at 1–15 days were at the highest risk ($P < 0.001$), and a significant relationship between season and infection ($P < 0.05$) was recorded. A significant association between infection and hygiene ($P < 0.001$), type of floor ($P < 0.01$) and source of water ($P < 0.01$) was also recorded. Statistical analysis concerning the clinical signs and fecal characteristics revealed a significant association with fecal consistency ($P < 0.001$), presence of blood ($P < 0.01$) and mucous ($P < 0.01$). Moreover, a significant association was found between infection and the desire for suckling ($P < 0.05$) and tenesmus ($P < 0.05$). The results of the present study demonstrated the strong relation between infections by *Cryptosporidium* spp. and diarrhea in buffalo calves.

45

Theriogenology

(2008) 70 : 1024–1029



Holding bovine oocytes in the absence of maturation inhibitors: Kinetics of in vitro maturation and effect on blastocyst development after in vitro fertilization

H. Alm a,* , Y.H. Choi b, L. Love b, B. Heleil c, H. Torner a,
K. Hinrichs b

a FBN Research Institute for the Biology of Farm Animals, 18196 Dummerstorf, Germany

b College of Veterinary Medicine and Biomedical Sciences, Texas A&M University, College Station, TX 77843, USA

c Faculty of Veterinary Medicine, Kafr El Sheikh University, Egypt

Received 7 March 2008; received in revised form 28 May 2008; accepted 28 May 2008

Abstract

Holding immature oocytes before maturation simplifies the transport of oocytes and aids in scheduling later manipulations. We examined the effect of holding bovine oocytes in the absence of meiotic inhibitors on their subsequent meiotic and developmental competence. Oocytes were matured immediately after recovery (control) or were held in a mixture of 40% TCM 199 with Earle's salts, 40% TCM 199 with Hanks' salts, and 20% FBS, at room temperature for 16 to 18 h (EH-held) and then matured. Chromatin status was determined at 0, 10, 14, 18, and 22 h of maturation culture. Oocytes were fertilized in vitro after either 18 or 22–24 h maturation. The EH treatment maintained oocytes at the germinal vesicle stage (79.3%, vs. 87.7% for control oocytes at 0 h; $P > 0.05$). Upon culture, held oocytes matured more quickly than did control oocytes. The proportions of mature oocytes were not significantly different between groups at 18 h (EH-held, 80.6% and control, 79.3%); however, after 22 h significantly more EH-held than control oocytes had degenerated (24.1% vs. 4.5%, $P < 0.0001$). Blastocyst development was similar between groups for oocytes fertilized after 18 h maturation (EH-held, 29.6% and control, 27.8%). When oocytes were fertilized after 22–24 h maturation, EH-held oocytes yielded lower blastocyst development than did control oocytes (16.5% vs. 29.3%, $P < 0.05$). In conclusion, bovine oocytes may be effectively held in the EH treatment before maturation without adversely affecting meiotic or developmental competence. However, holding affects the kinetics of maturation and this must be taken into account when subsequent manipulations are performed.