

المادة: ديناميكا تحليلية

الفرقة: الثالثة

الزمن: ساعتين

التاريخ: ١٧-١٨-٢٠١٨



جامعة كفر الشيخ

كلية العلوم

قسم الرياضيات

Answer the following questions

Q(1)

(a) Find the components of velocity and acceleration of a point in cylindrical polar Coordinates.

(b) A spherical pendulum is started horizontally at the level of the center. Prove that the equations of motion in azimuth and depth are

$$(i) (l^2 - z^2)\dot{\phi} = V l \quad (ii) l^2 \ddot{z} = z [2g(V^2 - \dot{z}^2) - V^2 \dot{z}]$$

Q(2)

(a) Find the components of velocity and acceleration for the motion of a particle on A twisted curve.

(b) A particle of mass  $m$  is projected horizontally under gravity with velocity  $v$  from a point on the inner surface of a smooth sphere of radius  $a$  at an angle distance  $\alpha$  from the lowest point find the pressure on the surface when it is at an angular distance  $\theta$  from the lowest point.

Q(3)

(a) Find the angular momentum of rigid body about a fixed point


(b) Find the equations of motion of a rigid body in three dimensions

Q(4)

Discuss the motion of a sphere on a rough plane under the action of forces the resultant of which passes through the center of the sphere

مع خالص تمنياتي بالتحقيق والنجاح

الاختبار النهائي الفصل الاول ٢٠١٧-٢٠١٨ م  
التاريخ: ٢٤ / ١٢ / ٢٠١٧ م  
الدرجة: ٧٠ درجة  
الزمن: ١٠-١٢



جامعة كفر الشيخ - كلية العلوم  
قسم الرياضيات  
الفرقة: الثالثة  
الشعبة: رياضيات  
المادة: توبولوجي

Solve the following questions:

Question (1)

(18 Marks)

Let  $\tau$  be the class of subsets of  $N$  consisting of  $\emptyset$  and all subsets of  $N$  of the form

$$E_n = \{n, n+1, n+2, \dots\} \text{ with } n \in N.$$

- (i) Show that  $\tau$  is a topology on  $N$ , (ii) list the open sets containing the positive integer 7.  
(ii) Find the family of closed subsets of  $(N, \tau)$ .

Question (2)

(18 Marks)

- a- If  $(X, D)$  is a discrete topological space and  $A$  is a subset of  $X$ , find  $A'$ .  
b- If  $(X, \tau)$  is a topological space and  $A, B$  are subsets of  $X$ , prove that  $(A \cap B)' = A' \cap B'$ .  
c- Show that a constant function is continuous.

Question (3)

(17 marks)

- a- If  $X = \{a, b, c\}$  and  $\beta$  is a family of subsets on  $X$  such that  $\beta = \{\{a, b\}, \{b, c\}\}$ , is  $\beta$  a base for any topology on  $X$ .  
b- Prove that a mapping  $f: (X, \tau) \rightarrow (Y, \sigma)$  is continuous if and only if  $f^{-1}(A) \subseteq (F^{-1}(A))$  for any  $A \subseteq Y$ .

Question (4)

(17 Marks)

- a- If  $(X, \tau)$  is a topological space and  $A$  is a subset of  $X$ , prove that  $A^- = A^{\circ} \cup A'$ .  
b- Prove that the intersection of two neighbourhoods of a point is also a neighbourhood of it.

End Questions

Best regards

Prof. Dr. Ahmed E.L. Maghrabi

س (3) (أ) أوجد قيمة التكامل  $\int \frac{zdz}{(z-4)^2(z^2+16)}$  باستخدام نظرية كوشي ونظرية

الباقي في الحالات التالية:

(ب)

(1)  $C : |z - 4i| = 1$  ..... (2)  $C : |z + 4i| = 1$

(3)  $C : |z - 4| = 1$  ..... (4)  $C : |z| = 1$  ..... (5)  $C : |z| = 6$

س (3) (أ) برهن نظرية لورانت.

(ب) أوجد متسلسلة لورانت التي تمثل الدالة  $f(z) = \frac{1}{(5-z)(z-6)}$  في المجالات الآتية:

(1)  $5 < |z| < 6$  ..... (2)  $|z| < 5$  ..... (3)  $|z - 6| < 1$



جامعة كفر الشيخ  
كلية العلوم

قسم الرياضيات امتحان الفصل الدراسي الأول للعام  
2017-----2018 م

الفرقة: المستوى الثالث (رياضيات)  
المادة: تحليل مركب (1) (320)  
الزمن: ساعتان  
التاريخ: الاربعاء 2017/12/27

اجب عن الأسئلة الآتية :

س (1) (أ) أوجد جذور المعادلة:  $Z^5 = -1 + \sqrt{3}i$

(ب) أدرس نقاط اتصال الدالة التالية:  $f(z) = \begin{cases} \frac{z^3 + i}{z - i} & , z \neq i \\ -6 & , z = i \end{cases}$

(ج) عرف الدالة التحليلية  $f(z)$  عند نقطة  $z_0$ .

(د) باستخدام معادلتى كوشي ريمان أي من الدوال تحليلية:

$$(1) f(z) = \frac{z}{z + (1 + 2i)} \quad (2) f(z) = ze^{2z}$$

(2) (أ) إذا كانت  $f(z) = u(x, y) + iv(x, y)$  وكانت المشتقات الجزئية  $u_x, u_y, v_x, v_y$

موجودة ومتصلة عند  $z_0$  فبرهن أن الدالة  $f(z)$  قابلة للاشتقاق عند  $z_0$  والمشتقة

$$f'(z_0) = u_x(x_0, y_0) + iv_x(x_0, y_0)$$

(ب) إذا كانت  $f(z)$  تحليلية على منطقة مترابطة  $D$  تحتوي على منحنى  $C$  مغلق بسيط موجب

$$\int_C \frac{f(s) ds}{(s - w)} = 2\pi i f(w) \dots, \notin C \text{ ان } w \text{ نقطة داخل } C \text{ فبرهن ان}$$

بقية الأسئلة بالخلف

(10 Marks)

Question 2

By the graph method find the optimal solution of the following (LPP):-

$$\text{Max } z = 2x_1 + 3x_2$$

$$\text{s.t. } x_1 + x_2 \leq 4, \quad 6x_1 + 2x_2 \geq 8, \quad x_1 + 5x_2 \geq 4$$

$$x_1 \leq 3, \quad x_2 \leq 3$$

$$\text{and } x_1, x_2 \geq 0$$

(20 Marks)

Question 3

Find the optimal solution of the following (LPP) and its duality

$$\text{Min } f(x) = 20x_1 + 16x_2$$

$$\text{s.t. } x_1 \geq 2.5, \quad x_2 \geq 6$$

$$2x_1 + x_2 \geq 17, \quad x_1 + x_2 \geq 12$$

$$\text{and } x_1, x_2 \geq 0$$

With Best Wishes Dr. Amin Elfeky



**Answer the following questions**

**Question1**

**(40 Marks)**

- 1- - State all types of duality linear programming problems ?
- 2- Prove that the intersection of any number of convex sets is also convex set?
- 3- Show that whether the following function is convex or concave :-

$$f(x) = x^2 + 2x$$

- 4- Prove that the dual of the dual is the primal ?
- 5- find the dual linear programming problem of the following :-

$$\text{Min } Z = x_1 + 4x_2 - 2x_3 - x_4$$

$$\text{s.t. } x_1 - 3x_2 + x_3 - x_4 \leq 2, \quad -x_1 + x_2 - 2x_3 = 3, \\ x_1 + 2x_2 - 3x_4 \geq 5, \quad -3x_2 - 2x_3 + x_4 \leq 6$$

And  $x_1, x_3 \geq 0$  ,  $x_2, x_4$  unrestricted variables

$\Rightarrow$  Look Page 2

**Answer the following questions**

**Question1**

(15 Marks)

Make use the following data, to estimate  $f''(2)$

X	0	1	3	4
Y	-3	-5	9	25

(25 Marks)

**Question2**

(15 Marks)

1- Prove that

$$i) \Delta^n f_{p-n} = \delta^n f_{p-\frac{n}{2}}, \quad ii) \nabla = -\frac{1}{2}\delta^2 + \delta\sqrt{1 + \frac{1}{4}\delta^2}$$

2- - if  $f(x) = u(x).v(x)$ , show that,

$$f[x_0, x_1] = u(x_0).v[x_0, x_1] + u[x_0, x_1].v(x_1)$$

(10 Marks)

(35 Marks)

**Question3**

(20 Marks)

1- Solve the following problem by two different methods :-

$$y' = x + y^2, \quad y(1) = -1, \quad h = 0.1, \quad \text{in } [1, 2]$$

compute the following

(15 Marks)

$$\int_1^4 \int_0^2 \frac{1}{(x+y)^2} dx dy, \quad n_x = 2, \quad n_y = 3$$

find the error.

With Best Wishes Dr. Amin Elfeky



Answer the following question

(1) (a) For any two events A, B prove that

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

(b) let A and B be events with  $P(A) = \frac{1}{2}$ ,  $P(B) = \frac{1}{8}$ ,  $P(A \cap B) = \frac{1}{4}$  find

(i)  $P(A \setminus B)$  (ii)  $P(B \setminus A)$  (iii)  $P(A \cup B)$  (iv)  $P(A^c \setminus B^c)$

(2) Given the probability density function

$$f(x) = \left. \begin{array}{ll} \frac{1}{6}x + k & 0 \leq x \leq 3, \\ 0 & \text{elsewhere} \end{array} \right\}$$

- (i) What is the value of k  
(ii) What is the cumulative distribution function of x  
(iii) Compute  $P(1 \leq x \leq 2)$ ,  $P(x \leq 3)$

(3) Team A has probability  $\frac{2}{5}$  of winning whenever it plays. If A plays 4 games, find the probability that A wins

- (i) 2 games (ii) at least one game (iii) more than half of the games.

(4) let x be a random variable with the standard normal distribution, find

- (i)  $P(0 \leq x \leq 1.42)$  (ii)  $P(-0.73 \leq x \leq 0)$   
(iii)  $P(-1.37 \leq x \leq 22.01)$  (iv)  $P(x \geq 1.13)$ .

(5) from the tables find mean, variance and standard deviation

class	2-8	9-15	16-22	23-29	30-36
f	5	8	6	3	2





Answer the following question

(1) Given the joint density function

$$f(x, y) = \begin{cases} e^{-\frac{x}{y}} & 0 < x < \infty, \quad 0 < y < \infty \\ 0 & \text{elsewhere} \end{cases}$$

(i) Find the marginal densities  $g(x)$ ,  $h(x)$

(ii) Find the conditional density  $f(x|y)$

(iii) Find  $E(x|y)$

(iv) Evaluate  $p(1 < x < 2 | y = 3)$

(2) Suppose that  $X$  and  $Y$  has the following joint probability function:

X \ Y	-3	2	4	Sum
1	0.1	0.2	0.2	0.5
3	0.3	0.1	0.1	0.5
sum	0.4	0.3	0.3	

(i) Find the marginal distributions of  $x$  and  $y$

(ii) Find  $\text{cov}(x, y)$

(iii) Find Correlation coefficient between  $X$  and  $Y$

(iv) Are  $x$  and  $y$  independent random variables?

(3) Let  $x$  and  $y$  are two independent random variables and let  $z$  be their sum. Find

(i) PMF of  $z$  when  $x$  and  $y$  are Binomial random variables.

(ii) PMF of  $z$  when  $x$  and  $y$  are Poisson random variables.

(4) Let  $x$  and  $y$  are independent and let  $w = x + y$ ,

$$f_x(x) = \lambda e^{-\lambda x}, x \geq 0,$$

$$f_y(y) = \lambda^2 y e^{-\lambda y}, y \geq 0$$

Find PMF of  $w$  and C.D.F.

الاختبار النهائي الفصل الأول ٢٠١٧ - ٢٠١٨ م  
اليوم: الأحد ١٤/١٨/٢٠١٨ م  
الزمن: ١٠-١٢  
الدرجة: ٧٠ درجة



جامعة كافر الشيخ - كلية العلوم  
قسم الرياضيات  
الفرقة: الثالثة (نبات - حيوان)  
المادة: رياضيات حيوية  
رمز المقرر: MATH 202

Solve the following questions:

Question (1)

( 18 Marks )

- a- Is the system of the linear equations a unique or infinite solution or no solution by Determinants:  
 $-3y + 2x + 5z = 3, 4x - y + z = 1, 3x - 2y + 3z = 4.$
- b- Find the final velocity of a car which is initially going at  $20 \text{ m/s}$  and then accelerates at a rate of  
 $a = 5 \text{ m/s}^2$  for a period of 15 seconds.
- c- If the following values represented the length of eight students by centimeters:  
 $173, 175, 168, 164, 172, 167, 170, 165.$  Find the median and the variance.

Question (2)

( 18 Marks )

- a- Derive the equation of the line passing through the point  $(-2, 3)$  and perpendicular to the line  
 $2x - 3y + 6 = 0.$
- b- By graphical method solve the simultaneous equations:  $y - x^2 = 0, y + 0.5x = 1.5.$
- c- A bacterial population initially with  $N_0 = 5.2 \times 10^5$  cells per m L is decaying exponentially according to  
 the equation  $N_t = N_0 e^{kt}$  and it is found that after  $t = 2$  hours the population has become  
 $N_t = 3.5 \times 10^4$  cells per m L, calculate the value of  $k$  in the equation.

Question (3)

( 17 Marks )

- a- Calculate the growth factor  $g$ , if (i) a population increase by 15% every ten years,  
 (ii) a bacterial population decrease by 25% every hour.
- b- If  $N_t = N_0 e^{-k(t-t_0)}$ , where  $N_t$  is the population at the time  $t$ ,  $N_0$  is the population at the time  
 $t = 0$ ,  $e$  is the fundamental exponential,  $k$  is the rate of growth of the population, find the value of  $t$

Question (4)

( 17 Marks )

- a- If  $N_t = N_0 e^{kt}$ , where  $N_t$  is the population at the time  $t$ ,  $N_0$  is the population at the time  
 $t = 0$ ,  $e$  is the fundamental exponential,  $k$  is the rate of growth of the population and  $t$  is the time.  
 Drive the linearization equation and hence find the slope and the intercept.
- b- Find the area of the circle with the center at  $(-4, 2)$  and tangent to the line  $3x + 4y - 16 = 0.$

End Questions

Best regards

Prof. Dr. Ahmed E.L. Maghlabi